

Original Article

Evaluation of Nanoemulsion of *Eucalyptus globulus* Oil as Potent Botanical Larvicide against Malaria Vector, *Anopheles stephensi* and West Nile Vector, *Culex pipiens* Under Laboratory and Semi-Field Conditions

Sara Sabzalizade¹, Amir Amani^{2,3}, Hassan Vatandoost^{1,4}, Fatemeh Hosseini¹,
Roya Najafi-Taher³, *Hamid Reza Basseri¹

¹Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

²Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran

³Medical Biomaterials Research Center (MBRC), Tehran University of Medical Sciences, Tehran, Iran

⁴Department of Chemical Pollutants and Pesticides, Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding author: Prof. Hamid Reza Basseri, email: basserih@tums.ac.ir

(Received 31 Aug 2021; accepted 08 Dec 2021)

Abstract

Background: Due to undesired environmental impact of insecticides as well as resistant of vectors to them, the development of organic and natural insecticides has been more considered. In the current study, we developed nanoemulsion of eucalyptus and investigated larvicidal activity of it against malaria vector, *Anopheles stephensi* and *Culex pipiens* under laboratory as well as semi-field conditions.

Methods: An optimized nanoemulsion was prepared by mixing Eucalyptus oil, Tween 80 and ethanol at ratio of 1:2:1.5 in distilled water, then, stirred for 20 minutes at room temperature. The product was then used for bioassay tests against 3-4th instar larvae of *Anopheles stephensi* as well as *Culex pipiens*. Furthermore, a semi-field trial was carried out to evaluate larvicidal activity of nanoemulsion of eucalyptus.

Results: Nanoemulsion of eucalyptus showed significantly high larvicidal activity comparing with bulk eucalyptus essential oil. The LC₅₀ and LC₉₀ value of nanoemulsion against *An. stephensi* were 111.0 and 180.8 ppm respectively and 29.5 and 73.7 ppm for *Cx. pipiens*, respectively. In the semi field condition, the Nanoemulsion of eucalyptus decreased 1-2nd instar larval density of Culicines and Anophelines to 90.1% and 85.2%, respectively.

Conclusion: The nano formulation of eucalyptus oil showed high larvicidal activity. Therefore, nanoemulsion of eucalyptus oil can be used as an eco-friendly larvicide against mosquitoes.

Keywords: *Eucalyptus*; Essential oil; Nanoemulsion; Larvicide; *Anopheles stephensi*;
Culex pipiens

Introduction

Mosquitoes are a serious threat to public health as they act as vectors that help in transmission of diseases that can be lethal (1,2). Owing to the lack of proper medication and

vaccines for treating mosquito-borne diseases, an alternative and effective approach used to control the vector population at the larvicidal stage is necessary because, during this stage,

the mosquitoes are in a stationary phase (3-5). The resistance to synthetic pesticides and harmful effects of their accumulation in the environment has created the need for natural and non-persistent insecticides. The essential oils extracted from plants are suitable as they are economically reasonable and have high activity in certain cases and are also biodegradable (4,6,7). Eucalyptus is a diverse genus of flowering trees and shrubs in the Myrtle family, Myrtaceae. The oil extracted from eucalyptus leaves possesses allelopathic property and prevents insects from attacking it, thereby, acting as a natural pesticide (8,9). The fumigation activity (10) and repellency (11) and insecticidal activity of eucalyptus oil has been demonstrated (12). Nanoemulsion of natural oils could enhance pesticide activity of the component (13). Nanoemulsions are emulsions whose droplet size is uniform and extremely small with the size ranging from 20 to 100 nm. Nanoemulsions are metastable and their stability is determined by the method of preparation (14). Nanoemulsions can be formulated by two kinds of methods such as high-energy and low-energy emulsification methods. The high-energy emulsification method comprises high-pressure homogenization and ultrasonication (15). Ultrasonication is the most widely used method owing to its ease of use and it is an economical method. The low-energy emulsification technique comprises methods that exploit the chemical properties of a system to convert a microemulsion into a nanoemulsion (16,17). The use of nanopesticides would be a contemporary measure for the control of pests and reducing the toxic effect of synthetic bulk pesticides on the environment (18). Recently, controlling for vector borne diseases using neem oil nanoemulsion will be of good alternative against *Culex quinquefasciatus* compared to the synthetic pesticides (13). The present study was carried out to develop a nano formulation of eucalyptus oil as an ecofriendly larvicide and evaluate its larvicidal activity against the larvae of malaria vector, *Anopheles stephensi* and vectors of West Nile virus (WNV), *Culex pipiens* in laboratory and semi-field conditions.

Material and Methods

Preparation of nanoemulsion

Eucalyptus oil (*Eucalyptus globulus*) was purchased from Barich Co., Iran and stored at room temperature under laboratory conditions; Tween 80 (Polyoxyethylene 20 monooleate) was supplied from Sigma. All other chemicals used were of analytical reagent grade. The oil-in-water nanoemulsion was formulated using eucalyptus oil, non-ionic surfactant (tween 80) and water. The concentration of eucalyptus oil (6%, v/v) was fixed for all the formulations. Initially, coarse emulsion was prepared by adding water to organic phase containing oil, surfactant and co-surfactant in ratios 1:1: 1.5 (v/v) using a magnetic stirrer, which was then subjected to ultrasonic emulsification using a 20 kHz Sonicator (Ultrasonics, USA) with a power output of 750 W. Energy input was given through sonotrode containing a piezoelectric crystal with a probe diameter of 13 mm. Sonicator probe generates disruptive forces that reduce the droplet diameter converting coarse emulsion to nanoemulsion. Then the characterization of nanoemulsions was carried out and the emulsion stability was investigated.

Droplet size distribution and polydispersity index

The droplet size distribution (analysis by volume) and poly dispersity index (PDI) of eucalyptus oil nanoemulsion formulation (1:2) was determined using a 90-plus particle size analyzer. The PDI is a measure of the homogeneity and stability of the droplet size in the nanoemulsion system. PDI values below 0.2 indicate a narrow size distribution and thus provide long-term stability to the formulated nanoemulsion. Prior to experiment, formulated emulsion was diluted with milli-Q (Millipore corporation) double-distilled water to trim down multiple scattering effects.

Morphology of emulsion droplets

To visualize the shape and morphology of

the formulated nanoemulsions, Atomic force microscopy (AFM) was carried out. One drop of emulsion was negatively stained with phosphotungstic acid and positioned on a copper grid.

Larvicidal bioassay

Mosquito rearing: Mosquitoes consisting of *An. stephensi* and *Cx. pipiens* were reared under a uniform condition including larvae and adult nutrition, temperature (28 ± 2 °C), humidity (70 ± 10 %) and on a 12-h light-dark cycle. Larvae were grown in bowls at a density of 200 larvae / 500 ml of distilled water with 0.01% table salt, and fed on fish food. The pupae were transferred to cages made of muslin cloth before eclosion to the adult stage. The female mosquitoes were fed on 10% fructose and the females were fed on guinea pig blood.

Larvicidal activity: The larvicidal activity of eucalyptus nanoemulsion against third instar larvae of *An. stephensi* and *Culex pipiens* were treated with different concentrations of nano and bulk eucalyptus oil emulsions. The bioassays were carried according to the guide line of World Health Organization for laboratory and field testing of mosquito larvicides (19). Initially, serial dilutions of nanoemulsion (5, 50, 100, 160, and 240 ppm) were prepared in sterile glass beakers (250 mL) containing 200 ml of water. Then 20 larvae (stage 3-4) of the mosquito species were placed into each beaker. The same procedure was conducted for the bulk of eucalyptus oil. The mortality rate of larvae was recorded after 24 and 48 hours. Each test was performed in four replicates. The percentages of larval mortality and standard deviation were calculated for each concentration of nanoemulsion and bulk emulsion.

Semi-field larval bioassays

Twelve artificial breeding places each 1×1 meters were provided in a semi-field condition in the Kazeroun area, southern Iran and allow the wild mosquitoes to lay their eggs on the surface of breeding places. The density of larvae 10 per dipper were

measured in each breeding place before any intervention. The spray of three times of obtained lethal concentration (LC₉₀) of bulk-eucalyptus oil (≈ 114 ppm) and nano-eucalyptus oil (≈ 90 ppm) were sprayed randomly on the surface of each breeding place. Each treatment was replicated in four breeding places and four breeding places of free oils as the control.

Statistical analysis

The mortality quantities of 50% and 90 % of imagicides (LC₅₀ and LC₉₀) and the level of confidence of 95%, the equation of the regression line were estimated using a regression probit analysis as described by Finney (20).

Mosquito identifications

All the adults females were identified using morphological identification key identification (21).

Results

Characterization of the selected nano-emulsion

Based on the thermodynamic stability study, 1:2 ratio nanoemulsions was selected, and their average diameter is 18 nm and their size distribution as presented in Fig.

1. The polydispersity index (PI) of the nanoemulsions is 0.060 which shows that it is uniform. The particles are presented in Atomic force microscopy (AFM) from 22 to 40 nm, and the droplets are spherical in nature (Fig. 2).

Larvicidal activity of eucalyptus oil nano-emulsion

The larvicidal activity of both nanoemulsion eucalyptus oil and bulk eucalyptus oil against larvae of *An. stephensi* and *Cx. pipiens* was varied. High mortality rate was observed among larvae of *An. stephensi* exposed to 160 ppm of nanoemulsion of eucalyptus oil within 24 hours, while the mortality rate for eucalyptus oil was 74% at the concentration of 160 ppm after 24 hours (Table 1). Moreover,

100% mortality occurred for the larvae exposed to both nanoemulsion eucalyptus oil and eucalyptus oil at a concentration of 240 ppm. Overall, the larvicidal activity of nanoemulsion oil against larvae of *An. stephensi* within 24 hours was significantly higher than eucalyptus at 160 ppm, but absolute mortality was observed among the larvae of *An. stephensi* after 48 hours in both nanoemulsion formulation and bulk oil of eucalyptus (Table 1). Similarity, the larvicidal activity of nanoemulsion oil against larvae of *Cx. pipiens* was higher than eucalyptus oil at 160 ppm within 24 hours while 100% mortality occurred by 240 ppm for both formulations of eucalyptus oils (Table 2). Lethal concentration (LC₅₀ and LC₉₀) for nanoemulsion of

eucalyptus oil and eucalyptus oil were significantly different within 24 and 48 hours after treatment (Table 3). Moreover, LC₅₀ and LC₉₀ of nanoemulsion of eucalyptus oil against larvae of *An. stephensi* significantly lower than eucalyptus oil within both 24 and 48 hours treatment (Table 3). Similarly,

lower LC₅₀ and LC₉₀ was observed for larvae of *Cx. pipense* exposed to nanoemulsion of eucalyptus oil which was significantly different with LC₅₀ and LC₉₀ for the larvae exposed to eucalyptus oil (Table 4).

Semi field larval bioassays

Both nano-eucalyptus and bulk eucalyptus oils decreased the density of mosquito larvae in the treated breeding places one day after




Fig. 1. Size distribution of eucalyptus nanoemulsion oil measured by Dynamic Light Scattering




Fig. 2. Atomic force microscopy image and size of eucalyptus nanoemulsion oil




Fig. 3. Residual larvicidal activity of Nanoemulsion of eucalyptus oil and bulk eucalyptus oil against different stages of mosquito larvae at semi-field condition. Twelve artificial breeding places (1×1 meters) were prepared and then allowed the wild mosquitoes laid eggs. L1, 2 = Larvae stage 1 & 2, L 3,4= Larvae stage 3 & 4

Table 1. Probit analysis of larvicidal activity of essential and nanoemulsion of *Eucalyptus globulus* oil on 3rd and 4th instar larvae of *Anopheles stephensi* after 24 hours

Concentration (ppm)	Essential oil			Nanoemulsion oil		
	Mortality (%)	Observed probit mortality	Expected probit mortality	Mortality (%)	Observed probit mortality	Expected probit mortality
60	9	3.659	3.246	8	3.595	3.384
160	74	5.643	5.658	91	5.025	5.204
240	100	7.526	6.654	100	7.576	7.024
360	100	7.526	7.651	100	7.576	8.089
Control	0	-	-	0	-	-

Table 2. Probit analysis of larvicidal activity of essential and nanoemulsion of *Eucalyptus globulus* oil on 3rd and 4th instar larvae of *Culex pipiens* after 24 hours

Concentration (ppm)	Essential oil			Nanoemulsion oil		
	Mortality (%)	Observed probit mortality	Expected probit mortality	Mortality (%)	Observed probit mortality	Expected probit mortality
60	8	3.595	3.384	11	3.773	3.520
160	51	5.025	5.204	86	4.900	5.161
240	100	7.576	7.024	100	7.576	6.802
360	100	7.576	8.089	100	7.576	8.443
Control	0	-	-	0	-	-

spraying. The high larvae population at stage 1 and 2 in the control breeding sites indicated that the wild mosquitoes laid eggs frequently during the bioassay period (Fig. 3). Therefore, the density of mosquito larvae was relatively high in the control breeding site while rapidly

declining in the treated sites. However, nano-eucalyptus emulsion kept the larvae density low for 6 days post spraying while the number of larvae at stage 1 and 2 increased two days after eucalyptus oil spraying (Fig. 3). The results indicating that nano-eucalyptus oil

Table 3. Lethal concentration (LC₅₀ and LC₉₀) and in the 24 and 48 hours bioassay tests of *Eucalyptus* oil and Nano-*Eucalyptus* against 3rd and 4th instar larvae of *Anopheles stephensi*

Exposure time (hours)	p-Value	χ^2 table (df)	χ^2 (Heterogeneity)	LC ₅₀ (ppm) \pm 95% C.L.	LC ₉₀ (ppm) \pm 95% C.L.	b \pm SE
<i>Eucalyptus</i> oil	24	0.01	15.086 (5)	103.7845 122.8343 145.0833	169.1937 206.2336 295.0632	5.6608 \pm 0.764
	48	0.00	15.035 (5)	93.761 101.663 145.0833	142.402 155.973 175.732	2.399 \pm 1.023
Nano- <i>Eucalyptus</i> oil	24	0.01	9.210 (2)	63.7924 80.8730 105.7719	102.8420 111.0171 119.7206	6.0457 \pm 0.539
	48	0.00	9.145 (2)	68.235 76.869 95.582	46.235 58.595 65.548	7.0457 \pm 0.652

Table 4. Lethal concentration (LC₅₀ and LC₉₀) and in the 24 and 48 hours bioassay tests of *Eucalyptus* oil and Nano-*Eucalyptus* against 3rd and 4th instar larvae of *Culex pipiens*

Exposure time (hours)	p-Value	χ^2 table (df)	χ^2 (Heterogeneity)	LC ₅₀ (ppm) \pm 95% C.L.	LC ₉₀ (ppm) \pm 95% C.L.	b \pm SE
<i>Eucalyptus</i> oil	24	0.01	8.235 (2)	47.3526 68.6321 76.1356	20.3120 39.4112 78.6321	6.5324 \pm 1.034
	48	0.01	9.210 (2)	42.4037 64.2180 74.0175	14.7673 37.3711 86.4146	.4508 \pm 1.127
Nano- <i>Eucalyptus</i> oil	24	0.01	9.210 (2)	43.3775 53.6733 70.0869	25.7675 29.4644 33.0781	3.2200 \pm 0.313
	48	0.01	9.451 (2)	31.6885 34.5681 39.0263	19.9645 23.2523 26.2879	5.1234 \pm 0.4123

has slightly longer residual larvicidal activity rather than bulk eucalyptus.

Discussion

Due to numerous limitations to control of adult mosquitoes, the ideal method to control them is targeting the larval stage. Control of mosquito larvae is based largely on the use of synthetic chemicals. Generally, synthetic pesticides have some disadvantages

such as health problems, harmful to the environment, pests may develop tolerance to certain chemicals over time and contamination of soil and water resources (22-24). However, the indiscriminate and injudicious use of pesticides has led to the widespread development of resistance among pests as well as insect vectors (25). Thus, alternative components to be needed for controlling mosquito vectors. Botanical pesticides are considered as safe, easily

biodegradable, environmentally friendly and with low toxicity (26,27). However, usage of them is often limited due to instability and rapid degradation, application frequency as well as requiring higher application rates. Hence, the use of botanical insecticides associated with nanotechnology offers considerable potential for increasing efficacy of plant based insecticide (27-29). Eucalyptus oil is the oil distilled from the leaves of eucalyptus, a genus of the plant family Myrtaceae. The repellent activity of eucalyptus oil was demonstrated against *Cx. quinquefasciatus* (30,31). The seed and leaf extract of eucalyptus oil contain compounds that are toxic to mosquito larvae (32,33). Since they do not cause any adverse health effects; they are used as insect repellents and a safe and eco-friendly alternative to synthetic pesticides (34). Therefore, in this study, we used eucalyptus oil as a safe and non-toxic larvicide. Generally, botanic pesticides have low effective durability; therefore, we utilized nanotechnology to overcome this problem. We synthesized nano-eucalyptus oil to increase larvicidal durability and effectiveness. The method of preparation determines the stability of the formulated nanoemulsion. Based on atomic force microscopy analysis, the size of our product was from 22 to 40 nm (Fig. 2), and the larvae mortality we achieved in laboratory and semi field bioassays (Tables 1, 2 and Fig. 3) was as the results of these nano-size. Nevertheless, the larvicidal effects of nano-eucalyptus oil are likely to change as the nano-particle size changes.

The stabilization of the nanoemulsions is also dependent on the steric effect of the non-ionic surfactant (35). Therefore, the effectiveness of nano-eucalyptus oil may be influenced by chemical component of water in the breeding sites. In semi-field assay, we did not observe 100% mortality of mosquito larvae a day after treatment of nano-eucalyptus oil and eucalyptus. This result may be due to components of water in the breeding sites. In addition, only a single concentration of nano-eucalyptus oil (≈ 90 ppm) was tested against wild mosquito

larvae. According to WHO guide line, the larvicide dosage for the field trial should be three times of LD₉₀ concentration in a laboratory scale. However, we did not observe convenient results for larvicidal effect of nano-eucalyptus oil in the field condition, Thus, more serial concentrations of the nano emulsion should be assays to overcome the highest larvae mortality in the field condition. In this study, eucalyptus oil (6%, v/v) is mixed with a non-ionic surfactant, tween 80 along with water, which is used as an aqueous phase. This emulsion is then subjected to ultrasonication that breaks down the bulk emulsion into an emulsion comprising droplets having the size in the nanometer range. This increases the surface area of the droplets, thereby, increasing the reactivity; thus, making nanoemulsions more effective than its bulk counterpart. However, the insecticidal and antimicrobial activity of eucalyptus oil is linked to its chemical composition such as 8-cineole (eucalyptol), which its concentration ratio varies in eucalyptus trees from different locations (36).

Conclusion

The nano formulation of eucalyptus oil showed high larvicidal activity against mosquito larvae when compared to its bulk counterpart. Therefore, nanoemulsion of eucalyptus oil can be used as an eco-friendly larvicide against mosquitoes. From this study, it can be concluded that eucalyptus oil nanoemulsion is a safe and effective alternative larvicide to control mosquito-borne diseases.

Acknowledgements

The authors would like to thank the School of Public Health, Tehran University of Medical Sciences, for their support in carrying out this research. This study was supported by the Deputy of Research, Tehran University of Medical Sciences, Grant No. 29565. The ethical approval was obtained from the ethics committee of Tehran

University of Medical Sciences.

Competing interests

The authors declare that they have no competing interests.

References

- Lajeunesse MJ, Avello DA, Behrmann MS, Buschbacher TJ, Carey K, Carroll J, Chafin TJ, Elkott F, Faust AM, Fauver H (2020) Infected mosquitoes have altered behavior to repellents: A systematic review and meta-analysis. *J Med Entomol.* 57:542-550.
- Nadim SS, Ghosh I, Martcheva M, Chattopadhyay J (2020) Impact of venereal transmission on the dynamics of vertically transmitted viral diseases among mosquitoes. *Math Biosci.* 325: 108366.
- Stoops CA, Qualls WA, Nguyen TT, Richards SL (2019) A review of studies evaluating insecticide barrier treatments for mosquito control from 1944 to 2018. *Environ Health Insights.* 13, 1178630219859004.
- Sukumar K, Perich MJ, Boobar LR (1991) Botanical derivatives in mosquito control: a review. *J Am Mosq Control Assoc.* 7: 210-237.
- Tamarina NA (1975) Natural control of blood-sucking mosquito population densities (review of literature). *Med Parazitol (Mosk).* 44: 603-607.
- Ochoa SA, Sanchez-Torres LE, Nevarez-Moorillon GV, Camacho AD, Noguera-Torres B (2017) Essential oils and their components as an alternative in the control of mosquito vectors of disease. *Biomedica.* 37:224-243.
- Fallatah SA, Khater EI (2010) Potential of medicinal plants in mosquito control. *J Egypt Soc Parasitol.* 40:1-26.
- Park JH, Lee HS (2018) Toxicities of *Eucalyptus dives* oil, 3-Carvomenthenone, and its analogues against stored-product insects. *J Food Prot.* 81: 653-658.
- Batish DR, PalSing H, Kumar Kohli R, Kaura S (2008) Eucalyptus essential oil as a natural pesticide. *For Ecol Manag.* 256: 2166-2174.
- Juan LW, Lucia A, Zerba EN, Harrand L, Marco M, Masuh HM (2011) Chemical composition and fumigant toxicity of the essential oils from 16 species of *Eucalyptus* against *Haematobia irritans* (Diptera: Muscidae) adults. *J Econ Entomol.* 104: 1087-1092.
- Mossi AJ, Astolfi V, Kubiak G, Lerin L, Zanella C, Toniazzo G, Oliveira D, Treichel H, Devilla IA, Cansian R, Restello R (2011) Insecticidal and repellency activity of essential oil of *Eucalyptus* sp. against *Sitophilus zeamais* Motschulsky (Coleoptera, Curculionidae). *J Sci Food Agric.* 91: 273-277.
- Alzogaray RA, Lucia A, Zerba EN, Masuh HM (2011) Insecticidal activity of essential oils from eleven *Eucalyptus* spp. and two hybrids: lethal and sublethal effects of their major components on *Blattella germanica*. *J Econ Entomol.* 104: 595-600.
- Anjali CH, Sharma Y, Mukherjee A, Chandrasekaran N (2012) Neem oil (*Azadirachta indica*) nanoemulsion--a potent larvicidal agent against *Culex quinquefasciatus*. *Pest Manag Sci.* 68: 158-163.
- Salvia-Trujillo L, Soliva-Fortuny R, Rojas-Grau MA, McClements DJ, Martin-Belloso O (2017) Edible Nanoemulsions as Carriers of Active Ingredients: A Review. *Annu Rev Food Sci Technol.* 8: 439-466.
- Kumar M, Bishnoi RS, Shukla AK, Jain CP (2019) Techniques for formulation of nanoemulsion drug delivery system: A review. *Prev Nutr Food Sci.* 24: 225-234.
- Sarheed O, Shouqair D, Ramesh K, Khaleel T, Amin M, Boateng J, Drechsler M (2020) Formation of stable nanoemulsions by ultrasound-assisted two-step emulsification process for topical drug delivery: Effect of oil phase composition and surfactant concentration and loratadine as ripening inhibitor. *Int J Pharm.* 576: 118952.
- Wang, L, Mutch KJ, Eastoe J, Heenan RK, Dong J (2008) Nanoemulsions prepared by a two-step low-energy process. *Langmuir.* 24: 6092-6099.
- Kumar S, Nehra M, Dilbaghi N, Marrazza G, Hassan AA, Kim KH (2019) Nano-based smart pesticide formulations: Emerging opportunities for agriculture. *J Control Release.* 294, 131-153.
- World Health Organization (2005) Guidelines for laboratory and field testing of mosquito larvicides. WHO/CDS/WHOPEP/GCDPP/2005.13. pp39
- Finney DJ (1971) Probit Analysis. 3rd ed. Cambridge University press, London. pp 333
- Azarian-Hamidi S, Harbach RE (2009). Keys to the adult females and fourth-instar larvae of the mosquitoes of Iran (Diptera: Culicidae). *Zootaxa.* 2078(1), 1–33.
- Yassin M, Ton J, Rolfe SA, Valentine TA, Cromei M, Holden N, Newton AC (2021) The rise, fall and resurrection of chemical-induced resistance agents. *Pest Manag Sci.* 10.1002/ps.6370
- Gurusubramanian G, Rahman A, Sarmah M, Ray S, Bora S (2008) Pesticide usage pattern in tea ecosystem, their retrospects and alternative measures. *J Environ Biol.* 29: 813-826.
- Casida JE (2016) Unexpected metabolic reactions and secondary targets of pesticide action. *J Agric*

- Food Chem. 64: 4471-4477.
25. Smith RF, Calvert DJ (1976) Health-related aspects of integrated pest management. *Environ Health Perspect.* 14: 185-191.
 26. Jaoko V, Nji Tizi Tanning C, Backx S, Mulatya J, Van den Abeele J, Magomere T, Olubayo F, Mangelinckx S, Werbroeck SPO, Smagghe G (2020) The phytochemical composition of *Melia volkensii* and its potential for insect pest management. *Plants (Basel).* 9(2): 143.
 27. Casida JE (2012) The greening of pesticide-environment interactions: some personal observations. *Environ Health Perspect.* 120: 487-493.
 28. Chaudhari AK, Singh VK, Kedia A, Das S, Dubey NK (2021) Essential oils and their bioactive compounds as eco-friendly novel green pesticides for management of storage insect pests: prospects and retrospects. *Environ Sci Pollut Res Int.* 28: 18918-18940.
 29. Luiz de Oliveira J, Ramos Campos EV, Fraceto LF (2018) Recent developments and challenges for nanoscale formulation of botanical pesticides for use in sustainable agriculture. *J Agric Food Chem.* 66: 8898-8913.
 30. Tian Y, Dong F, Zhou X, Yang X (2020) Repellent, insecticidal and antimicrobial activities of leaf essential oils from three eucalyptus species. *Chem Biodivers.* 17:e1900580.
 31. Magesa SM, Kamugisha ML (2006) Evaluation of the bio-efficacy of three brands of repellents against wild populations of anthropophilic mosquitoes. *Tanzan Health Res Bull.* 8: 145-148.
 32. Cheng SS, Huang CG, Chen YJ, Yu JJ, Chen WJ, Chang ST (2009) Chemical compositions and larvicidal activities of leaf essential oils from two eucalyptus species. *Bioresour Technol.* 100: 452-456.
 33. Senthil Nathan S (2007) The use of *Eucalyptus tereticornis* Sm. (Myrtaceae) oil (leaf extract) as a natural larvicidal agent against the malaria vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Bioresour Technol.* 98: 1856-1860.
 34. Kweka EJ, Moshia F, Lowassa A, Mahande AM, Kitau J, Matowo J, Mahande MJ, Massenga CP, Tenu F, Feston E, Lyatuu E, Mboya M, Mndeme R, Chuwa G, Temu E (2008) Ethnobotanical study of some of mosquito repellent plants in north-eastern Tanzania. *Malar J.* 7:152.
 35. Perugini L, Cinelli G, Cofelice M, Ceglie A, Lopez F, Cuomo F (2018) Effect of the coexistence of sodium caseinate and Tween 20 as stabilizers of food emulsions at acidic pH. *Colloids Surf B Biointerfaces.* 168: 163-168.
 36. Gonzalez A, Gutierrez-Cutino M, Moenne A (2014) Oligo-carrageenan kappa-induced reducing redox status and increase in TRR/TRX activities promote activation and reprogramming of terpenoid metabolism in Eucalyptus trees. *Molecules.* 19: 7356-7367.