

Letter to the Editor

The Tick of the Museum: New Record of *Amblyomma helvolum* on *Naja sumatrana*

*Dagmara Dyczko¹, Bartosz Borczyk²

¹Division of Biological Sciences, Department of Microbial Ecology and Acaroentomology, University of Wrocław, Wrocław, Poland

²Division of Biological Sciences, Department of Evolutionary Biology and Conservation of Vertebrates, University of Wrocław, Wrocław, Poland

*Corresponding author: Dr Dagmara Dyczko, E-mail: dagmara.dyczko2@uwr.edu.pl

(Received 05 Mar 2026; accepted 23 June 2026)

Dear Editor

We report a historical record of the reptile-associated tick *Amblyomma helvolum* Koch, 1844 parasitizing the equatorial spitting cobra *Naja sumatrana* Laurenti, 1768, based on a preserved specimen housed in the herpetological collection of the Naturhistorisches Museum Wien (Natural History Museum Vienna, NMW), Austria.

The tick *A. helvolum* is an Asiatic species typically inhabiting tropical and subtropical forest ecosystems throughout Southeast Asia (1, 2). It is primarily a reptile-associated tick, commonly parasitizing snakes and lizards (3–6). However, a few exceptional records indicate that this species may occasionally feed on birds and mammals, including humans (7, 8). Although parasitism of *N. sumatrana* by this tick has been reported previously (6, 9–11), available records are rare and fragmentary, likely reflecting the nocturnal habits and high venom toxicity of the host, which limit opportunities for direct observation and examination. In this context, the present record contributes to the existing knowledge by providing a museum-vouchered historical occurrence of *A. helvolum* on *N. sumatrana* from Sumatra, together with information on the precise attachment site.

Re-examination of zoological museum specimens can provide valuable insights into his-

torical host-parasite relationships, revealing parasitic associations and distribution patterns that are often inaccessible through contemporary field studies, including those involving reptile-associated ticks (12–14). Museum-vouchered parasite records are especially valuable because they provide independently verifiable evidence of host association, locality and collection date, thereby strengthening historical datasets used to reconstruct parasite distribution and host use.

The observation was made incidentally during a morphological examination of preserved snake specimens conducted for purposes unrelated to parasitological research. Among 33 examined specimens of *N. sumatrana* (all available specimens in the NMW collection), two ticks were found on a single male specimen (catalogue no. NMW 27809-5), collected in Padang (Sumatra) in July 1896. The ticks were attached to the ventral surface of the tail. They were identified as adult females of *A. helvolum* (Fig. 1) based on visible external diagnostic characters and published species descriptions (8, 15). As the ticks were not removed from the host specimen, their identification was based on photographic documentation and non-invasive assessment of externally visible morphological characters. The characters considered included an ovoid idiosoma, the presence of

...

<https://doi.org/10.18502/jad.v20i2.22411>

festoons, a pair of very subtle dorsal foveae below the scutum, and an ornate scutum that is wider than long, with three distinct pale ornamented patches: one anterolateral to each eye and one at the posteromedial margin of the scutum. Because the ticks remain attached to the historical host specimen, no microscopic examination, measurements, or destructive sampling were performed. The ticks are retained in situ on the host specimen *N. sumatrana* NMW 27809-5 in the herpetological collection of the Natural History Museum Vienna and have not been assigned a separate catalogue number. This record thus adds a precisely documented nineteenth-century museum-based occurrence to the limited historical evidence for the association between *A. helvolum* and *N. sumatrana*.

Ticks were detected on only one of 33 examined *N. sumatrana* specimens, suggesting that this species may represent an occasional host for *A. helvolum*. Although limited, such observations provide valuable context for understanding host–parasite interactions and aspects of the natural history of both the snake and the tick.

A review of published host records by Oda et al. (6) documented 113 records of *A. helvolum*, of which only five originated from Sumatra, including a single record from *N. sumatrana*. All known records from this island are historical and exceed 75 years in age, including the present specimen collected in 1896. The lack of more recent records may reflect limited knowledge of tick distribution in the region, low abundance of the species in natural habitats, or changes in its distribution and abundance.

Ticks parasitizing snakes are most commonly found on the head and neck region, whereas attachment to the body is considered uncommon (4, 5). The observation of ticks on the snake's tail is therefore unusual. The host specimen exhibited tail damage, although its timing and origin cannot be determined from the preserved material. Therefore, any asso-

ciation between this damage and tick attachment should be interpreted with caution. A similar observation was reported by Lazell et al. (16), who documented *A. helvolum* attaching to sites around an old wound on the posterior part of the body of the Sulawesi black racer *Ptyas dipsas*. The unusual tail attachment observed here therefore represents an additional biological detail that was not evident from previous host lists or broader reviews and may contribute to future assessments of attachment-site selection in reptile hosts.

This record, based on a single historical specimen, adds to the limited knowledge of host associations of *A. helvolum* and underscores the value of museum collections as archives of past host-parasite interactions, particularly for rare or hazardous host species that are difficult to study under field conditions. More specifically, it complements previous syntheses by providing a dated, museum-vouchered Sumatran record from *N. sumatrana*, together with information on an unusual attachment site.

...
<https://doi.org/10.18502/jad.v20i2.22411>



Fig. 1. *Amblyomma helvolum* ticks attached to the ventral surface of the tail of *Naja sumatrana*, preserved specimen (catalogue no. NMW 27809-5), collected in Padang (Sumatra) in July 1896, housed in the herpetological collection of the Naturhistorisches Museum Wien (Natural History Museum Vienna, NMW), Austria

Acknowledgements

We would like to thank Silke Schweiger and Georg Gassner (Natural History Museum Vienna, NMW, Austria) for granting access to the herpetological collection under their care. The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

References

1. Kolonin GV (1995) Review of the ixodid tick fauna (Acari: Ixodidae) of Vietnam. *J Med Entomol.* 32: 276–282.
2. Petney TN, Keirans JE (1995) Ticks of the genera *Amblyomma* and *Hyalomma* from South-East Asia. *Trop Biomed.* 12: 45–56.
3. Amarga AKS, Lin JW, Tseng HY, Robbins RG, Kwak ML, Lin SM (2023) New host records of the reptile tick *Amblyomma helvolum* Koch (Ixodida: Ixodidae: Amblyomminae) in Taiwan. *Syst Appl Acarol.* 28: 931–943.
4. Chao LL, Hsieh CK, Shih CM (2013) First report of *Amblyomma helvolum* (Acari: Ixodidae) from the Taiwan stink snake *Elaphe carinata* (Reptilia: Colubridae), collected in southern Taiwan. *Ticks Tick Borne Dis.* 4: 246–250.
5. Mao JJ, Norval G, Robbins RG, Wong ST (2019) An instance of *Boiga dendrophila* (Boie, 1827) (Reptilia: Colubridae) being parasitized by *Amblyomma helvolum* Koch, 1844 (Acari: Ixodidae), with comments about the attachment sites of this tick species. *Acarologia.* 59: 115–119.
6. Oda FH, Martins TF, Labruna MB, O'Shea M, Kaiser H (2023) Ticks (Acari: Ixodidae) of three Timor-Leste reptiles: first country record of *Amblyomma helvolum*, with new interactions and an updated list of host species. *Ticks Tick Borne Dis.* 14: 102060.
7. Guglielmone AA, Robbins RG, Apanaskevich DA, Petney TA, Estrada-Peña A, Horak IG (2014) *The hard ticks of the world.* Springer, Dordrecht.
8. Kwak ML, Jones MD, Harman MEA, Smith SN, D'Souza A, Knierim T, Barnes CH, Waengsothorn S, Amarga AKS, Kuo C-

...

<https://doi.org/10.18502/jad.v20i2.22411>

- C, Nakao R (2023) Taxonomy, biology and new host records of the East Indies reptile tick *Amblyomma helvolum* Koch, 1844 (Acari: Ixodidae), including the first record of human infestation. Ticks Tick Borne Dis. 14: 102224.
9. Anastos G (1950) The scutate ticks, or Ixodidae of Indonesia. Entomol Am. 30: 1–144.
 10. Kohls GM (1957) Malaysian Parasites-XVIII. Ticks (Ixodoidea) of Borneo and Malaya. Stud Inst Med Res Fed Malay States. 28: 65–94.
 11. Kho KL, Koh FX, Tay ST (2015) Molecular evidence of potential novel spotted fever group rickettsiae, *Anaplasma* and *Ehrlichia* species in *Amblyomma* ticks parasitizing wild snakes. Parasit Vectors. 8: 112.
 12. Arzúa M, Onofrio V, Barros-Battesti D (2005) Catalogue of the tick collection (Acari, Ixodida) of the Museu de História Natural Capão da Imbuia, Curitiba, Paraná, Brazil. Rev Bras Zool. 22: 623–632.
 13. De Jesus C, Bhosale C, Wilson K, White Z, Wisely S (2021) Reptile host associations of *Ixodes scapularis* in Florida and implications for *Borrelia* spp. Ecology. Pathogens. 10: 999.
 14. Wood CL, Vanhove MPM (2023) Is the world wormier than it used to be? We'll never know without natural history collections. J Anim Ecol. 92: 250–262.
 15. Voltzit OV, Keirans JE (2002) A review of Asian *Amblyomma* species (Acari, Ixodida, Ixodidae). Acarina. 10: 95–136.
 16. Lazell JD, Keirans JE, Samuelson GA (1991) The Sulawesi black racer, *Coluber* (*Ptyas*) *dipsas* and a remarkable ectoparasite aggregation. Pac Sci. 45: 355–361.